CALLUS INDUCTION AND SHOOT REGENERATION FROM SESAME SEEDS

Osisami-Onibiyo Olubukunola F[1]*,Ajenifujah-Solebo S.O[1],Adeboye Seyi E[1], Udofia E. G[1], Hassan Abu S[1], Adeshina D.A[1]

1Plant Tissue Culture, Agricultural Biotechnology Department, National Biotechnology Development Agency, NABDA, Umaru Musa Yar'adua Expressway, Abuja, Nigeria

Corresponding Author: Email: osisamiolubukunola@yahoo.com Tel: 08064891720

Abstract

In this work, the responsiveness of several explants of sesame seedlings (*Sesamum indicum L.*) to the induction of callus and shoot formation is assessed. Sesame is one of the oldest and most out-of-date oilseed crops grown for its edible oil, which is utilized in food preparation and other industrial uses. The seeds are rich in nutrients and have many positive health effects. However, Nigeria's output potential has not yet been completely harnessed. The goal of this study was to create a method for in vitro callus induction (CI) and shoot regeneration from sesame seedling explants. For the in vitro callus induction (CI) and shoot regeneration of the plantlets, the explants were cultured in a base medium containing Murashige and Skoog (MS) supplemented with various concentration ratios of auxins and cytokinines. After the second week of inoculation, callus induction (CI) started to develop. When compared to all other combinations, Kinetin and 2, 4-dichlorophenoxyacetic acid (2, 4-D) were more successful for callus induction (CI) frequencies in all explants, while MS with 0.5ms BAP and 0.75mls NAA showed the greatest number of shoot regenerations.

Keywords: Callus Induction (CI), in vitro, sesame, shoot regeneration

Introduction

Sesame (*Sesamum indicum L.*), often known as benne, is a flowering plant in the genus Sesamum. Sesame has been grown for many years in India, Pakistan, Burma, China, Japan, Indo-China, and Africa. Sesame has only recently been brought to Mexico, Central America, South America, and the United States (Lalpantluangi and Shah, 2018). It is one of Ethiopia's main oilseed crops. Sesame is thought to have originated in Ethiopia, where there is a large genetic diversity that can be used as a resource to further enhance the crop (Daniel and Parzies, 2011).

Sesame seeds are full of vital elements like folic acid, protein, unsaturated fatty acids, oil, and vitamins and minerals. It was domesticated more than 3,000 years ago, making it one of the oldest oilseed crops ever. Of all seeds, it contains the most oil. Ethiopia is regarded as the center of origin for sesame, and the genetic variety is significant, acting as resources for further crop improvement, while India ranks first in both area and production and also has the leading position for export of white seeded type, which are in great demand (Daniel and Parzies, 2011).

It is regarded as the oldest oil seed crop known to man and is known as the queen of oil seed crops. It is one of the most significant oil seed crops and has been grown in tropical regions since ancient times (Weiss, 1971). However, conventional breeding only makes small genetic improvements. The goal of the current work is to create procedures for the callus induction and shoot regeneration of seedlings from various explants (Were et al., 2006).

The in vitro callus induction and shoot regeneration of explants were examined in this work (Younghee, 2001). For the purpose of growing plants from the cotyledon, secondary leaves, stem, and roots explants of Sesamum indicum L., the effects of various hormones on the callus, the start of various explants in MS media, and the reproducible methodology were also described.

Materials and Methods

Preparation of explants

Sesamum indicum L. seeds were purchased for experimental work from a well-known market (Nyanya) in Abuja, Nigeria, and transported to the National Biotechnology Development Agency, NABDA, Lugbe, Abuja, Nigeria for additional study at the Tissue Culture Laboratory, Plant Improvement Unit of the Agricultural Biotechnology Department. The sesame seeds were rinsed for 20 minutes with tap water before being steeped in a 3% Tween 20 solution for five minutes. The soaked seeds were repeatedly rinsed with sterile distilled water. The seeds were then thoroughly cleaned again with sterile distilled water after being disinfected for 45 minutes at room temperature with 70% ethanol. The final step in the surface sterilization process for explants was a thorough three-time washing with sterile distilled water after a further five minutes of surface cleaning with a 2% solution of sodium hypochlorite. The explants were then inoculated into 2.5 x 15 cm test tubes filled with non-absorbent cotton. After that, the cultures spent 14 days in a light photoperiod. Explants were made from the 14-day aseptic seedlings' portions of cotyledon, secondary leaves, stem, and roots (Asad et al., 2019).

Preparation of media

The trials that followed employed MS basal medium supplemented with 3% (w/v) sucrose and 0.8% (w/v) agar. Depending on what was needed, HCl or NaOH was used

to bring the pH level down to 5.8. In order to prevent clumping, Phytagel (agar) was added to the medium at a rate of 3g/L slowly and gradually. The medium was then heated to create a homogenous solution. It was put in culture tubes after the agar medium had been added. The medium was sterilized after preparation in an autoclave at 121°C and 15 pounds of pressure for 15 minutes.

In 150 × 25 mm test tubes containing 10 ml of culture medium, segments of cotyledon, secondary leaves, stem, and roots explants were inoculated into the medium before being tightly covered with non-absorbent cotton. Explant cultures were stored in the growth room, which was lit by fluorescent tubes all the time. For purer cultures, sub-culturing was carried out at intervals of 20 days or longer (Asad et al., 2019)

In vitro shoot regeneration from sesame seeds

Cotyledon, secondary leaves, stem and root explants from in vitro-cultured seedlings that were 7 days old were introduced in a shoot regeneration medium containing MS salts (Murashige & Skoog, 1962), sucrose, vitamins (Gamborg et al., 1968), and various concentrations of BAP (0.5mls) and NAA (0.75mls), both individually and collectively (Ahmed et al., 2008)

In vitro callus induction

Explants of 7-day-old seedlings cultured in vitro were placed in a callus initiation medium containing MS salts, sucrose, and various concentrations of 2,4-D and NAA (0.4mls) alone or in combination with Kinetin (0.4mls) for the induction of callus. Cotyledon, secondary leaf, stem, and root segments were used as explants (Baskaran et al., 2006)

Result and Discussion

After the first week of inoculation of the four explants (cotyledon, secondary leaves, stem, and roots), the greatest number and maximum lengths of shoots were noticed (Seo et al., 2007). As can be seen in Table 1, the cotyledon and root explants exhibit the largest percentage of shoots in comparison to the other types. The stem explant gave no reaction in any of the media that were employed. After 3 weeks of inoculation, the four explants (cotyledon, secondary leaves, stem, and roots) showed signs of a friable, green, proliferating callus. Table 2's Cotyledon and Stem explants exhibit the greatest percentage of callus compared to the other explants on MS supplemented with 2, 4-D (2mg/l), and kinetin (0.5mg/l), compared to other combinations (Martin, 2002).

The secondary leaf and root explants on the same media did not respond (Table 2). Successful callus maintenance was attained for at least two cycles in the current experiment. According to the results, cotyledon and root explants responded well for shoot regeneration while cotyledon and stem explants responded well for the induction of calluses, therefore they are used as model explants for this purpose, as indicated in the table 2 below.

Conclusion

Plant tissue culture, sometimes referred to as in vitro micropropagation of plants, has become more significant in recent years due to its use in the quick multiplication, propagation, and ex situ preservation of particular plants. According to the findings of this study, Callus induction and plant regeneration in Sesame (*Sesamum indicum L.*) through direct seeding, (Lokesha et al., 2007), cotyledon and stem can be used as efficient explants for callus induction, whereas cotyledon and root are efficient

Table 1: Response (in per cent) of different explants of sesame seeds for shoot regeneration with varied concentrations of Plant Growth Regulators, NAA and BAP

		BAP	NAA (mg/l)	C	2^0L	S	R	Remarks of Shoots
Medium A, Week	**1**	0.5	-	10.0	15.0	0.0	10.0	Small viable shoot
	2			15.0	20.0	0.0	25.0	Good viable shoots
	3			30.0	30.0	0.0	35.0	Good viable shoots
	4			45.0	35.0	0.0	45.0	Good viable shoots
Medium B, Week	**1**	0.75	0.75	10.0	10.0	0.0	15.0	Small viable shoots
	2			20.0	15.0	0.0	20.0	Viable shoots
	3			25.0	25.0	0.0	25.0	Good viable shoots
	4			40.0	40.0	0.0	40.0	Good viable shoots
Medium C, Week	**1**	-	1	0.0	0.0	0.0	0.0	No visible shoot
	2			0.0	0.0	0.0	0.0	No visible shoots
	3			0.0	0.0	0.0	0.0	No visible shoots
	4			0.0	0.0	0.0	0.0	No visible shoots

Table 2: Response (in per cent) of different explant of sesame seeds for callus induction with varied concentrations of Plant Growth Regulators, 2, 4 D, NAA and Kinetin

MS +SUCROSE+	2,4 D (mg/l)	NAA	KN	C	2°L	S	R	Remarks of Callus
Medium A, Week 1	0.4	-	-	0.0	10.0	15.0	0.0	Little, greenish
2				0.0	25.0	35.0	0.0	Little, greenish
3				0.0	30.0	40.0	0.0	Little, greenish, friable
4				0.0	55.0	55.0	0.0	Greenish, globular good Callus
Medium B, Week 1	-	0.4	-	5.0	10.0	15.0	0.0	Little, greenish
2				15.0	20.0	25.0	0.0	Little, greenish
3				0.0	35.0	0.0	0.0	Littlegreenish, compact
4				0.0	60.0	0.0	0.0	Greenish, globular, compact good Callus
Medium C, Week 1	-	-	0.4	10.0	10.0	10.0	0.	Little, greenish
2				20.0	25.0	25.0	0.	Little, greenish friable
3				35.0	30.0	30.0	0.0	Little, greenish, compac
4				45.0	40.0	45.0	0.	Greenish,, globular compact good Callus
Medium D, Week 1	0.4	0.4	-	10.0	10.0	10.0	0.	Little, greenish
2				20.0	25.0	25.0	0.	Little, greenish friable
3				35.0	30.0	30.0	0.	Little, greenish, compac
4				45.0	40.0	45.0	0.	Greenish, globular good Callus
Medium E, Week 1	0.4	-	0.4	15.0	0.0	15.0	0.0	Little, greenish
2				30.0	0.0	30.0	0.0	Little, greenish, friable
3				45.0	0.0	45.0	0.0	Little, greenish, compact
4				55.0	0.0	55.0	0.0	Greenish, globular compact, good Callus

Note: C= Cotyledon, 2oL = Secondary leaves, S=Stem, R=Root



A B C

Figure 1: Different in vitro developmental stages of Sesamum indicum L
A- initiation of shoot from seeds; B- initiation of callus; C- callus induction maintenance (subcultured)

explants for seed-based shoot regeneration (Noman et al., 2008). It may be conceivable to regenerate an entire plant from genetically variable callus cells because callus culture is a good source of genetic or karyotype variety.

Recommendation

Further research on sesame seeds should be carried out.

Declaration of interest

The authors have no relevant financial or non-financial interests to disclose.

Acknowledgement

I sincerely thank the Director General, Prof .Abdullahi Mustapha for this wonderful opportunity, for creating NJBR-NABDA Journal of Biotechnology Research, which is the first of its kind in NABDA and also, Dr. Ajenifujah-Solebo O, Deputy Director, Agricultural Biotechnology Department, National Biotechnology Development Agency (NABDA) Lugbe, Abuja, my mentor and diligent boss, who took her time to supervise the work and still guiding me towards the successful completion of the work.

REFERENCES

1. Abdellatef, E., Ahmed, M. M, Daffalla, H. M. and Khalafalla, M. M. (2010). Enhancement of adventitious shoot regeneration in sesame (Sesamum indicum L.) cultivar Promoky using ethylene inhibitors. Journal of Phytomedicine 2:61-7
2. Ahmed, M. M., Abdellatef, E. and Khalfallah, M. M. (2008). In vitro multiple shoot induction and plant regeneration in elite Sudanese sesame cultivars (Sesamum indicum L.). Am Eurasian Journal of Sustainable Agriculture, 2:308-14.
3. Asad, M., Ahmed, N., Sohail, A., Sher, J., Burni, T. and Hadi, F., 2019. 32. In vitro callus induction and plantlet regeneration of sesame (Sesamum Indicum L.). Pure and Applied Biology (PAB), 8(2), 1307-1313.
4. Baskaran, P. and Jayabalan, N. (2006). In vitro mass propagation and diverses callus orientation on (Sesamum indicum L) an important Oil. Journal of Agricultural Technology, 259-69
5. Daniel E. G. and Parzies, H. K. (2011). Genetic variability among landraces of sesame in Ethiopia. African Crop Science Journal, 19(1):1-13
6. Gamborg OL, Miller RA & Ojima K (1968). Nutrient requirements of suspension culture of soybean root cells. Experimental Cell Research 50: 151-158
7. Janowicz J & Wojciechowski A (2009). The evaluation of regeneration ability of two flax (Linum usitatissimum L.) Cultivars in in vitro culture. Oilseed Crops 30(1): 35-50.
8. Lalpantluangi, P.C. and Shah, P., 2018. Character association and path coefficient analysis in Sesame (Sesamum indicum L.) genotypes under foothill condition of Nagaland. The Pharma Innovation, 7(5b):82
9. Lokesha R, Shashidhara N & Janagoudar BS (2007). Callus induction and plant regeneration in Sesame (Sesamum indicum L.) through direct seeding. Plant Cell Biotechnology and Molecular Biology; 8 (1&2): 85-88.
10. Martin KP (2002). Rapid propagation of Holostemma ada-kodien Schult, a rare medicinal plant, through axillary bud multiplication and indirect organogenesis. Plant Cell Reports 21: 112-117.
11. Murashige T & Skoog F (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. Plant Physiology 15:473-497
12. Noman, A.S.M., Islam, M. S., Siddique, N. A., Hossain, H. (2008). High frequency induction of multiple shoots from nodal explant of Vitex negundo using silver nitrate. International Journal of Agricultural Biology; 10:6
13. Seo HY, Kim YJ, Park TI, Kim HS, Yun SJ, Park KH, Oh MK, Choi MY, Paik CH, Lee YS & Choi YE (2007). High frequency plant regeneration via adventitious shoot formation from deembryonated cotyledon explants of Sesamum indicum L. In Vitro Cellular & Developmental Biology 43: 209-214
14. Were, B. G., Onkware, A. O., Carlsson, A. S. and Welander, W. (2006). In vitro regeneration of sesame (Sesamum indicum L.) from seedling cotyledon and hypocotyly explants. Plant Cell Tissue organ Culture; 85:235-9; DOI: 10.1007/s11240-006-9077-3
15. Weiss, E.A. (1971). Castor and Sesame and Safflower. Leonard Hill, London, 311-525.
16. Younghee K (2001). Effects of BA, AA, 2, 4-D and AgNO3 treatments on the callus induction and shoot regeneration from hypocotyl and cotyledon of sesame (Sesamum indicum L.). Korean Journal of Horticultural Science & Technology 42: 70-74.