

BDC/PS/003/2026**THE NIGER DELTA BIORESOURCES: THE NEED FOR AN INTEGRATED
PROGRAM FOR SUSTAINABLE EXPLOITATION AND CONSERVATION.****¹OHIMAIN, E. I., AND ²WOOD, T. T.****¹Department of Microbiology, Niger Delta University, Wilberforce Island,
Bayelsa State, Nigeria****²Bioresources Development Centre, Odi Bayelsa State****Correspondence: elijahohimain@ndu.edu.ng, 08037306520****Abstract**

The Niger Delta is a globally recognized biodiversity hotspot and one of the largest wetland ecosystems in the world, supporting diverse habitats such as mangroves, freshwater swamps, and tropical rainforests. These ecosystems host a significant array of flora and fauna, including diverse species of fish, which sustain the livelihoods of millions of people through ecosystem services like food security and raw material provision. However, decades of oil and gas exploration have led to severe environmental degradation, characterized by frequent oil spills, gas flaring, and dredging, which have resulted in habitat fragmentation and chronic pollution. Over-extraction of other bioresources including timber, fuelwood and non-timber forest products is impacting the ecosystem. Additionally, the spread of invasive species like water hyacinth and Nipa palm, which is actively displacing native mangroves, further threatening the region's ecological integrity. Current conservation efforts are often undermined by weak regulatory enforcement, lack of stakeholder coordination, and a failure to integrate local community interests. This paper proposes the development of an integrated management framework that reconciles biodiversity exploitation with ecological preservation through strategic policy formulation and multi-stakeholder engagement models. By identifying priority conservation strategies, such as the designation of protected areas and habitat restoration, the study provides a roadmap for the sustainable exploitation of the Niger Delta's bioresources while ensuring their long-term protection for future generations.

Introduction

The Niger Delta, located at the southern tip of Nigeria, is one of the largest wetland ecosystems in the world and is recognized as a critical global biodiversity hotspot (Angaye & Lelei, 2025; Ewim et al., 2023). This vast region encompasses various ecological zones, including mangroves, freshwater swamps, and tropical rainforests, which collectively host a remarkable array of flora and fauna (Ewim et al., 2023). Notably, the Delta's aquatic ecosystems contain over 200 species of fish, representing the highest concentration of fish biodiversity in any coastal ecosystem in West Africa (Chidumeje et al., 2015). Nwosu (2010) documented A total of 339 plant species belonging to 88 families have been identified in the forests and homestead gardens of the Niger Delta. These bioresources serve as the bedrock of the local economy, providing essential ecosystem services such as food security, medicinal resources, and raw materials including timber, fuel, and fibre. These bioresources supports millions of people who depend on forestry, fishing and agriculture for their survival (Adekola &

Mitchell, 2011; Ewim et al., 2023). These resources had sustained the region for centuries, before petroleum was discovered in the 1950s, whose exploitation has changed the economics and impacted the ecosystem of the region.

Despite its immense ecological and socio-economic value, the Niger Delta has been subjected to intensive resource exploitation, primarily driven by the oil and gas industry since the mid-20th century (Adekola & Mitchell, 2011; Angaye & Lelei, 2025). The environmental consequences of these activities have been profound. Dredging operations widen and deepen waterways in order to permit marine navigation. This drastically alters natural hydrology, increases coastal and riverbank erosion, and destroys delicate spawning grounds for aquatic species (Ohimain et al., 2003). Decades of frequent oil spills, continuous gas flaring, and extensive dredging have resulted in chronic soil and water pollution, which has decimated local fish populations and degraded agricultural land (Raimi, 2020; Ugochukwu & Ertel, 2012; Ohimain, 2022). Furthermore, the ecological balance is being further destabilized by the proliferation of invasive species, particularly nipa palm (*Nypa Fruticans*) and water hyacinth (*Eichhornia crassipes*). For instance, Nipa palm is aggressively displacing native mangrove forests, which are vital for coastal protection and carbon sequestration (Nwobi et al., 2020; Ogbeibu & Oribhabor, 2023), while water hyacinths are clogging the waterways and causing siltation rapid siltation (Aghoghovwia, & Obomunu, 2022; Onyari et al., 2024). Unsustainable exploitation of natural resources is also exacerbating natural disasters such as flooding, which affects biodiversity including wildlife and vegetation and threatens human survival (Ohimain et al. 2014a, 2014b).

The management of these bioresources is currently at a crossroads. While the region is central to Nigeria's national economy, the regulatory framework governing environmental protection remains largely ineffective (Onu, 2003). Issues such as weak policy implementation, inadequate monitoring, and a lack of transparency in the regulatory process have allowed environmental degradation to persist with minimal accountability (Moffat & Lindén, 1995; Ugochukwu & Ertel, 2012). Furthermore, the unsustainable exploitation of bioresources is mounting pressures on the ecosystem (Ohimain et al 2014a). There is an urgent need to shift from a focus on short-term resource extraction to a model of sustainable development that prioritizes biodiversity conservation and the well-being of local communities (Angaye & Lelei, 2025; Moffat & Lindén, 1995). The primary goal of this study is to provide a comprehensive review of the Niger Delta's bioresources and to propose a sustainable management framework.

Niger Delta Bioresources

The Niger Delta serves as a critical biological hub for Nigeria, characterized by a vast network of wetlands that host a significant percentage of the nation's species. These bioresources provide essential ecological functions and are the primary source of survival for the region's rural populations. Its environment encompasses four distinct ecological zones (UNDP, 2016), namely, coastal barrier islands, mangrove swamp forests, freshwater swamps, and lowland rainforests, forming a vastly well-endowed ecosystem that contains one of the Earth's highest concentrations of biodiversity.

The Niger Delta is home to a complex mosaic of habitats, including the largest mangrove forest in Africa and the third largest in the world (Sakib, 2021). These habitats are vital to Nigeria's overall biodiversity, as the Niger Delta is estimated to hold between

60% and 80% of the country's endemic plant and animal species (Sakib, 2021). The region's ecosystems ranging from saline mangroves to permanent freshwater swamp forests, support a portion of the national wildlife diversity, which includes approximately 7,895 plant species and over 22,000 vertebrate and invertebrate species (Izah, 2018). Among the notable fauna residing in these sensitive ecosystems are the mona monkey, speckle-throated otter, and various species of barracuda and catfish (Sakib, 2021).




Figure 1: Ecological zones of the Niger Delta (Source: UNDP, 2016)

The Niger Delta is endowed with diverse bioresources of various economic and environmental importance including plant species (Table 1), animal species (Table 2), microbes (Table 3), and even waste biomass, when utilized (Table 4). The plant species in the Niger Delta have been put to various social and economic uses including timber for building construction and furniture, wild fruits and food substances, fuelwood, traditional medicine, cultural uses and totems. The fauna has been similarly been used for food, construction material, totems, and pets. In addition, the region is blessed with fertile soil and good agricultural climate, which has been widely exploited for crop farming and aquaculture. Exploitation of biodiversity resources also results to the generation of various waste stream, which are often not treated but dumped. Thus, polluting the environment. Oil exploration has also resulted to spills that pollute the environment. Fortunately, there exists diverse species of microbiota that have been demonstrated to degrade wastes including oil spills.

The Niger Delta has great economic potential. The region's bioresources are central to the Nigerian economy, supporting both industrial sectors and local subsistence livelihoods. Some of the resources of the Nigeria Delta beyond petroleum that has been tapped include fisheries, agriculture and forestry. Economic sectors that exploit the Niger Delta bioresources include tourism, bioenergy, traditional medicine, timber and other building/construction materials. Fisheries and aquaculture are key economic activities in the region. The Niger Delta's coastal zones are highly productive, with more than 70% of the fish stocks targeted by industrial fisheries being caught in these waters (Chidumeje et al., 2015). Artisanal fisherfolks rely heavily on these resources, particularly small pelagic species which constitute over 50% of the total fish catch (Chidumeje et al., 2015). Mangroves of the Niger Delta are breeding and spawning grounds for fishes of the south Atlantic Ocean (Ogbeibu & Oribhabor, 2023; Yetunde, 2017).

Agriculture and Forestry are another important economic activity in the Niger Delta. The fertile soil and climate support a variety of cash crops, including palm oil, rubber, and cocoa (Ewim et al., 2023). Additionally, the wetlands provide essential timber and non-timber products such as saw logs, bamboo, raffia, and medicinal species (Nwankwoala & Okujagu, 2021). The partial microbial degradation of biomass in sedimentary basin is linked to the formation of petroleum hydrocarbons ab-initio (Head et al., 2003). Hydrocarbon extraction remains the dominant economic driver, with crude oil production from the region accounting for over 90% of Nigeria's total export earnings (Udotong et al., 2017). Ecosystem Services are key to survival in any environment. Beyond direct production, the Niger Delta ecosystem and associated biota provides regulatory services including flood control, water purification, and carbon sequestration, all of which are vital for regional environmental stability (Ewim et al., 2023).

Table 1: Selected floral resources of the Niger Delta

Bioresource	Species	Documented Uses	Reference
Mangrove Zone Species	<i>Rhizophora racemosa</i> , <i>R. mangle</i> , <i>R. harizonii</i> , <i>Avicennia germinans</i> , <i>Laguncularia racemose</i> , <i>Conocarpus erectus</i>	Construction poles, furniture and firewood; bark ash used for salt extraction; fish smoking, construction of fishing gear, leaf extracts used as mosquito larvicides against <i>Anopheles gambiae</i> , antimicrobial activity used for traditional medicine; stabilising coastal mudflats	Numbere 2018; Angaye et al. 2014, Baba et al. 2013; Ohimain et al., 2026
Freshwater Swamp Forest Species	<i>Elaeis guineensis</i> , <i>Raphia hookeri</i> , <i>Cocos nucifera</i> , <i>Mitragyna</i>	Fruits, oils, biofuel, fuelwood, timber and construction materials, traditional medicine, craft making	Ubom, 2010; Igu and Marchant, 2018; Diyaolu et al. 2022

	<i>ciliate, Hevea brasiliensis Irvingia gabonensis, Dacryodes edulis</i>		
<i>Lowland Rainforest Species</i>	<i>Khaya grandifoliola, Lophira alata, Ocimum gratissimum, Musanga cecropioides</i>	Timber, construction materials, furniture, traditional medicine, wild fruits,	<i>Diyaolu et al. 2022; Ubom 2010</i>
<i>Aquatic plants</i>	<i>Pistia stratiotes, Alternanthera nodiflora, Polygonum senegalense</i>	Traditional medicine, animal fodder, compost, phytoremediation of pollutants	<i>Ita 1994</i>

Table 2: Selected Fauna Resources of the Niger Delta

Bioresource	Scientific Name	Documented Uses	Reference
Mammals	<i>Piliocolobus epieni, Cercopithecus sclateri, Cercopithecus erythrogaster, Pan troglodytes ellioti, Loxodonta cyclotis, Hippopotamus amphibius, Civettictis civetta, Trichechus senegalensis, Smutsia gigantea, Syncerus caffer nanus, Philantomba maxwellii</i>	Conservation importance, bushmeat, seed disperser and forest regeneration agent; ecotourism value; cultural totem species, pets, traditional medicine, traditional festivals,	<i>Akani et al. 2015; Luiselli et al. 2015</i>
Reptile	<i>Crocodylus niloticus, Osteolaemus tetraspis, Python sebae Bitis gabonica, Varanus niloticus Pelusios niger, Pelomedusa subrufa, Mecistops cataphractus</i>	Skin used for luxury leather goods, totem, bushmeat; traditional medicine, conservation importance, craft	<i>Akani & Luiselli 2001; Akani et al. 2015</i>

Bioresource	Scientific Name	Documented Uses	Reference
Birds (Avian Fauna)	<i>Psittacus Erithacus</i> , <i>Ardea alba</i> , <i>Lophoceros semifasciatus</i> , <i>Haliaeetus vocifer</i> , <i>Megaceryle maxima</i> , <i>Bubulcus ibis</i> , <i>Actophilornis africanus</i>	Conservation importance, pet, cultural importance, seed disperser, traditional medicine, ecotourism and birdwatching, bushmeat	<i>Adeola 1992</i>
Fish (Ichthyofauna)	<i>Chrysichthys nigrodigitatus</i> , <i>Clarias gariepinus</i> , <i>Ethmalosa fimbriata</i> , <i>Heterotis niloticus</i> , <i>Tilapia guineensis</i> , <i>Mugil cephalus</i> , <i>Lutjanus endecacanthus</i> , <i>Pseudotolithus typus</i>	Source of protein, ecotourism, creates employment,	<i>Nwosu et al. 2021;</i>
Invertebrates (Aquatic and Terrestrial)	<i>Crassostrea gasar</i> , <i>Tympanotonus fuscatus</i> , <i>Callinectes pallidus</i> , <i>Penaeus notialis</i> , <i>Archachatina marginata</i>	Protein source, construction material, calcium supplements, decorative material, traditional medicine, cultural use, creates employment	<i>Egonmwan 2007, 2008; Nwosu et al. 2021</i>
Amphibians	<i>Hoplobatrachus occipitalis</i> , <i>Hyperolius sp</i> , <i>Amietophrynus regularis</i>	Bushmeat, traditional medicine, ecological role as pest control, part of food web, pets	<i>Akani et al. 2004</i>

Table 3: Microbial Resources of the Niger Delta

Microbial Resource	Species	Application
Hydrocarbon-degraders	<i>Pseudochrobactrum</i> sp., <i>Bacillus albus</i> , <i>Pseudomonas aeruginosa</i> , <i>Alcaligenes faecalis</i> , <i>Alcaligenes ammonioxydans</i> , <i>Cupriavidus malaysiensis</i> , <i>Proteus</i>	Bioremediation of oil spills, soil clean-up

	<i>vulgaris</i> , <i>Serratia</i> , <i>Micrococcus</i> , <i>Klebsiella</i> , <i>Corynebacterium</i> , <i>Rhodococcus</i> , <i>Alcanivorax</i> , <i>Arthrobacter</i> , <i>Alcaligenes</i> ammonioxydans, <i>Aspergillus</i> , <i>Penicillium</i> , <i>Rhizopus</i> , <i>Mucor</i> , and <i>Candida</i> .	
Photosynthesizing microbes	<i>Beggiatoa</i> , <i>Chloronema</i> , <i>Chromatium</i> , <i>Thiopedia</i> , <i>Leucothiobacteria</i> and several species of microalgae	Useful for primary productivity, biofuel production
Biosurfactant producers	<i>Pseudomonas aeruginosa</i> , <i>Alcaligenes</i> ammonioxydans and <i>Bacillus subtilis</i>	Oil dispersion, Microbial Enhanced Oil Recovery (MEOR), industrial detergents
Nitrogen fixing and nitrifying bacteria	<i>Azospirillum</i> , <i>Azotobacter</i> , <i>Rhizobium</i> , <i>Clostridium</i> , <i>Nitrobacter</i> sp., <i>Nitrosomonas</i> sp., <i>Klebsiella</i> <i>pneumoniae</i> , <i>Enterobacter mori</i> , and <i>Roseomonas rhizosphaerae</i> .	Help in plant nutrition and some species are also hydrocarbon degraders
Methanogens	<i>Methanococcus methylutens</i> ,	Synthesis and degradation of methane. Climate change relevance
Phosphate-solubilizers	<i>Bacillus</i> , <i>Pseudomonas</i> , <i>Enterobacter</i> , <i>Xanthobacter</i> , <i>Kluyvera</i> , <i>Paenibacillus</i> , <i>Chryseomonas</i> , <i>Vibrio proteolyticus</i>	Support plant nutrition
Sulphate reducers	<i>Desulfovibrio</i> , <i>Desulfococcus</i> , <i>Desulfotomaculum</i> , <i>Desulfosarcina</i>	Anaerobic hydrocarbon degradation. Microbial corrosion. Heavy metal bioremediation

Table 4: Biomass waste as a resource

Waste Type	Source in Delta	Est. Quantity	Current Use	Potential
Cassava peels	Garri/fufu processors in Rivers, Delta, Bayelsa	5-8 million tons/yr	Dumped, burnt, feed	H ₂ , biogas, ethanol, animal feed. 60-70% carbs
Palm Oil Mill Effluent POME	500+ palm oil mills	2.5-3.5 million m ³ /yr	Lagoons, river discharge	Biogas, H ₂ , fertilizer. High COD 25,000-75,000 mg/L

Palm kernel shell	Palm oil mills	1.2 million tons/yr	Boiler fuel, dumped	Briquettes, concrete
Empty fruit bunches EFB	Palm oil mills	1.8 million tons/yr	Boiler fuel, dumped	Cellulose for ethanol, compost, gasification
Oil palm fronds/trunks	Plantations	6-8 million tons/yr	Left in field	Pellets, biogas, pyrolysis oil
Rice husk/straw	Delta, Bayelsa rice farms	300,000 tons/yr	Open burning	Silica, briquettes, syngas
Coconut husk/shell	Coastal areas	150,000 tons/yr	Charcoal, dumped	Activated carbon, briquettes
Plantain/banana peels	Markets, households	400,000 tons/yr	Dumped	Biogas, H ₂ , compost. High starch
Corn wastes	Markets, households	4.1 million tons/yr for Nigeria	Dumped	Biogas, H ₂ , compost. High starch
Municipal solid wastes	Households	1.8-2.2 million tons/yr	Dumped	Highest H ₂ yield: 150-260 mL H ₂ /g VS; biogas, compost
Sewage/Greywater	Households	20 - 120 Liters/person/day	Dumped	Biogas, H ₂ , nutrient recovery
Abattoir waste	Slaughterhouses	50,000 tons/yr	Dumped	Biogas – high N, fats

Exploitation Challenges

The exploitation of the region's natural resources including fisheries, timber, non-timber forest products and hydrocarbon resources has created a legacy of environmental and socio-economic challenges. Besides, the generation of diverse wastes that are poorly managed including municipal solid wastes, agricultural processing wastes and oil and gas exploration wastes has exacerbated the environmental challenges of the region. While the region is a high conservation priority, it remains virtually unprotected, allowing industrial activities to severely disrupt its ecological integrity (Anejionu et al., 2014; Angaye & Lelei, 2025).

Unsustainable biodiversity extraction in the Niger Delta involves both industrial-scale resource exploitation and local artisanal over-extraction. Non-hydrocarbon exploration practices that have profound effects on biodiversity includes waste dumping on land and rivers, dredging and sand mining, agricultural expansion into virgin forests, overfishing and fishing with chemicals, aquaculture, fuel wood harvesting and picking of non-timber forest products including wild fruits such as palms, mango and bush mango, wine tapping, bush meat and other wildlife hunting such as snails, tortoises and turtles (Ohimain et al 2014a, 2014b; Ohimain, 2022). The most destructive practices include oil spills, gas flaring, destructive artisanal refining, overharvesting of mangrove timber, and unregulated overfishing (Angaye & Lelei, 2025).

Resource exploitation by indigenous people in the Niger Delta that has profound effects on biodiversity include illegal oil bunkering and artisanal refining, destructive fishing methods, wood logging and fuelwood gathering, solid waste dumping and bushmeat hunting (Angaye & Lelei, 2025; Onuoha & Uzodiana, 2026). Destructive fishing methods is a major source of impact on fisheries caused by local fisherfolks. Unregulated fishing pressures and the use of illegal gear such as fine-meshed nets, chemical poisons, and explosives can deplete fish stocks and kill non-target aquatic species indiscriminately (Angaye & Lelei, 2025). Mangrove and freshwater forests are aggressively cleared for urban settlement and cut down for fuelwood by local communities. This strips the region of its primary natural barrier against erosion and destroys the nursery ecosystems for countless species (Angaye & Lelei, 2025). Commercial bushmeat hunting is causing biodiversity impacts. The unregulated hunting of wildlife (especially primates and other mammals) for the bushmeat trade is a major threat to terrestrial biodiversity (Ohimain, 2022). The illegal bunkering and small-scale refining of crude oil in swamps and forested areas generates high-temperature fires and releases heavy, toxic crude oil residues directly into water bodies and onto adjacent vegetation (Angaye & Lelei, 2025).

Sand mining and wetland reclamation are also mounting pressure on the biodiversity of the Niger Delta. Sand extraction for civil construction purposes is quite common. The draining and filling of wetlands for agricultural expansion and urban real estate completely erase highly localized and sensitive freshwater and saltwater ecosystems (Onuoha & Uzodiana, 2026). The use of municipal wastes to reclaim wetlands is exacerbating the impacts on biodiversity and present a route for introduction of plastics into the aquatic system. Invasive species particularly nipa palm and water hyacinth are most problematic in the Niger Delta. The introduction and unchecked spread of nipa palm are fast displacing indigenous mangrove vegetation (Nwobi et al., 2020), while water hyacinth is clogging the waterways and causing rapid siltation (Aghoghovwia, & Obomunu, 2022; Onyari et al., 2024).

Environmental Degradation associated with oil exploration

The most immediate threat to the Niger Delta's bioresources is the chronic discharge of crude oil and the impact of gas flaring. Decades of leakages, equipment failure, and pipeline vandalism release millions of barrels of crude oil. Oil spill coats mangroves and swamps, decimating flora, destroying fish habitats, and introducing heavy metals into the soil and food chain (Onuoha & Uzodiana, 2026). For instance, in 2004, over 74,900 barrels were lost to the environment, while 2006 saw a massive loss of

approximately 428,196 barrels (Ude et al., 2023). More recent records highlight the ongoing nature of this crisis. For instance, Jonathan et al (2025) reported over 21,914 barrels of crude oil spill from the period 2012 to 2022, with only about 38% recovered. Illegal crude oil bunkering and artisanal refineries has exacerbated oil spills in the region, many of which go on unreported (Chris & Oghenetekevwe, 2023; Richard et al., 2022)

These spills have profound consequences for the ecosystem and biodiversity:

- **Habitat and Biodiversity Loss:** Oil spills and gas flaring are major drivers of biodiversity loss, placing numerous species of flora and fauna at heightened risk of extinction (Angaye & Lelei, 2025). Hydrocarbon contamination in the Bomu area, for instance, resulted in fires that devastated creeks and mangrove forests, leaving vast agricultural lands barren for decades (Ewim et al., 2023).
- **Water and Soil Pollution:** The "wide consequences" of extraction include the degradation of soil and water supplies, which directly affects human health as well as the survival of aquatic and terrestrial life (Angaye & Lelei, 2025).
- **Agricultural Impact:** Gas flaring and pollution have been linked to significant negative effects on crop farming, further threatening the food security of local communities (Ude et al., 2023).

The ecological integrity of the Niger Delta is currently under severe threat, primarily due to the "wide consequences of oil extraction," which include habitat deterioration and chronic pollution (Angaye & Lelei, 2025). Major ecological zones, specifically the lowland rainforests and barrier island forests, are reportedly heavily impacted (Sakib, 2021). This degradation has led to a noticeable decline in wildlife populations; several species once common to the region, such as the elephant, chimpanzee, and yellow-backed duiker, are now considered endangered or are facing extinction, with some species no longer being sighted in their traditional habitats (Izah et al., 2018). Oil spills and gas flaring have further compromised the environment by degrading soil and water supplies, which directly impacts the health of both local communities and the surrounding flora and fauna (Angaye & Lelei, 2025).

Concerns over Unsustainable Extraction Practices

There are risks posed by environmental degradation and unsustainable extraction practices on both the ecosystem and dependent human populations (Adekola & Mitchell, 2011; Ugochukwu & Ertel, 2012). The unsustainable extraction of natural resources is putting pressure on and altering the ecosystem of the Niger Delta and could be responsible for increasing natural disasters such as flooding (Ohimain et al., 2014b). The challenges of exploitation are compounded by practices that prioritize resource extraction over environmental sustainability. Concerns include;

- **Underreporting and Regulatory Gaps:** A critical issue is the lack of transparency in environmental monitoring; many oil spills are not officially reported (Chidumeje et al., 2015). Non-petroleum impacts are seldom reported for regulatory oversight.
- **Industrial Discharge and Agricultural Impacts:** Beyond oil spills, the discharge of industrial wastewater and agricultural runoff contributes to the overall pollution

of the Niger Delta's diverse habitats, including its freshwater swamps and tropical rainforests (Ewim et al., 2023).

- **Disruption of Livelihoods:** The deterioration of habitat and the pollution of water bodies disrupt traditional livelihoods, such as fishing and farming, which are the primary survival mechanisms for the local population (Angaye & Lelei, 2025).

Addressing these challenges requires a shift toward eco-friendly exploration technologies and the rigorous application of environmental standards to secure the well-being of future generations (Angaye & Lelei, 2025).

Integrated Management Framework

The survival of the Niger Delta's ecosystems requires a transition from fragmented, reactive management to an Integrated Management Framework (IMF) that harmonizes economic goals with ecological preservation. Current systems often fail due to a lack of coordination and the prioritization of short-term industrial gains over long-term environmental health (Moffat & Lindén, 1995; Ugochukwu & Ertel, 2012). An effective IMF serves as a roadmap for reconciling the intensive resource extraction typical of the region with the "urgent need for biodiversity conservation" (Angaye & Lelei, 2025). Integrated management framework that aligns strategic policy formulation with effective stakeholder engagement models involving government, industry, and local communities (Onu, 2003; Raimi, 2020) are urgent in order to preserve the Niger Delta biodiversity. Also, identification and recommendation of specific conservation strategies, such as the designation of new protected areas and large-scale habitat restoration initiatives, to mitigate the impacts of industrial activity and invasive species are required (Angaye & Lelei, 2025; Ogbeibu & Oribhabor, 2023). The integrated management framework for Niger Delta bioresources could involve policy formulation, stakeholders' engagement, conservation strategies, protected area designation and habitat restoration.

Strategic policy formulation is vital for sustainable extraction of resources in the Niger Delta. The foundation of an integrated framework lies in robust and transparent policy formulation. Currently, the Nigerian regulatory process suffers from performance gaps that hinder environmental protection (Onu, 2003). Strategic policies must move beyond symbolic regulation toward "result-oriented action plans" (Ogbeibu & Oribhabor, 2023). Key policy requirements include:

- **Strengthening Regulatory Independence:** Enhancing the autonomy and monitoring capabilities of environmental agencies to ensure they can hold oil and gas operators accountable for "chronic oil spills" and other violations (Onu, 2003; Ugochukwu & Ertel, 2012).
- **Enforcement of Environmental Standards:** Rigorous application of existing laws, such as those governing the impacts of dredging on marine biodiversity, which have historically been poorly managed (Raimi, 2020).
- **Integrating Sustainable Development Goals:** Ensuring that national economic policies for the Niger Delta prioritize the long-term viability of its bioresources alongside hydrocarbon production (Angaye & Lelei, 2025; Moffat & Lindén, 1995).

- **Regulatory oversight for artisanal resource extraction is required:** Most legal framework focuses on multinationals and other industrial activities, neglecting the numerous activities of local people.

Stakeholder engagement is key for integrated resource extraction strategy. Successful management is impossible without the active participation of all parties involved in the Niger Delta's landscape. A multi-stakeholder engagement model is necessary to bridge the gap between industry, government, and the people most affected by degradation (Onu, 2003; Raimi, 2020). Models for improved stakeholder involvement include;

- **Community-Based Management:** Empowering local communities by involving them in the decision-making process. This fosters a sense of ownership and utilizes indigenous knowledge for more effective biodiversity management (Angaye & Lelei, 2025; Ugochukwu & Ertel, 2012). The Akassa Sea Turtle Conservation Group and Bonny Environmental Committee are good examples.
- **Corporate Accountability:** Developing models that require oil companies to engage in transparent dialogue and collaborative restoration projects with host communities, moving beyond simple compensation to active partnership (Moffat & Lindén, 1995; Onu, 2003). The Hydrocarbon Restoration Project (HYPREP) is carrying out environmental restoration and mangrove restoration in Ogoni land.
- **Conflict Resolution:** Implementing frameworks that address the social tensions arising from environmental damage, thereby creating a stable environment where conservation strategies can be successfully applied (Onu, 2003).

By aligning these strategic and collaborative elements, the Niger Delta can move toward a model of development that preserves its status as a global biodiversity hotspot while maintaining its economic contributions (Angaye & Lelei, 2025; Moffat & Lindén, 1995).

Conservation Strategies and Habitat restoration

Conservation Strategies are key to the survival of the ecosystem. Given the "urgent need for biodiversity conservation" in the Niger Delta, implementing robust conservation strategies is critical to halting the rapid loss of species and habitat (Angaye & Lelei, 2025). Despite its global significance, the region remains virtually unprotected, necessitating a shift toward proactive environmental management (Anejionu et al., 2014).

Declaration of protected area status for the Niger Delta is key for their preservation. One of the most pressing challenges in the Niger Delta is that high-priority biodiversity areas currently lack formal protection from industrial and other anthropogenic encroachment (Anejionu et al., 2014; Angaye & Lelei, 2025). For instances, several forest reserves in the region have been encroached (Ohimain, 2022). Establishing a network of protected areas is essential for safeguarding the Delta's unique bioresources.

- **Targeting Sensitive Ecosystems:** Conservation efforts must prioritize the designation of legally protected zones in sensitive habitats, such as the lowland rainforests and barrier island forests, which are currently declining due to exploitation (Sakib, 2021; Ugochukwu & Ertel, 2012).

- **Mangrove Preservation:** As the region hosts the largest mangrove forest in Africa, specific "result-oriented action plans" are required to protect these coastal buffers from further fragmentation caused by oil infrastructure and dredging (Ohimain, 2004; Ohimain et al., 2005; Ogbeibu & Oribhabor, 2023; Raimi, 2020).
- **Representative Conservation:** Protected areas should be selected based on biodiversity patterns to ensure they cover the habitats of the region's 200+ fish species and endangered wildlife, such as the mona monkey and various endemic plants (Chidumeje et al., 2015; Sakib, 2021).

Habitat restoration initiatives are important in the integrated management of Niger Delta bioresources. Restoration is a vital secondary strategy to recover ecosystems that have already been degraded by decades of oil spills, industrial effluents and other anthropogenic activities (Angaye & Lelei, 2025).

- **Remediation of Contaminated Sites:** Active restoration is needed for oil spill impacted areas like the such as Bomu region, where hydrocarbon contamination has left vast tracts of mangrove forests and agricultural land barren for years (Ewim et al., 2023). Restoration must involve the removal of pollutants and the replanting of native vegetation to restore soil and water health (Angaye & Lelei, 2025).
- **Invasive Species Management:** A critical component of restoration involves the removal of invasive species. For instance, the Nipa palm is aggressively replacing native mangroves; restoration projects must focus on eradicating nipa palm and re-establishing native mangrove species to maintain coastal integrity (Ogbeibu & Oribhabor, 2023).
- **Community-Led Restoration:** Incorporating local communities into restoration projects ensures long-term success. By involving residents in habitat recovery, conservation strategies can simultaneously provide alternative livelihoods and leverage indigenous knowledge to improve ecosystem resilience (Moffat & Lindén, 1995; Ugochukwu & Ertel, 2012).

Through the combination of formal protected status and active ecological restoration, the Niger Delta can begin to mitigate the long-term impacts of resource extraction and secure its bioresources for future generations.

Conclusion

The Niger Delta stands as one of the world's most significant ecological treasures, characterized by a complex network of mangroves, freshwater swamps, and rainforests that support a vast array of biodiversity, including over 200 species of fish. However, this study underscores that the region's bioresources are currently in a state of crisis. Decades of intensive resource extraction including fisheries, timber and fuel wood, non-forest products and petroleum exploration have impacted the bioresources of the region. Hydrocarbon exploitation have led to the release of millions of barrels of oil into the environment, resulting in "wide consequences" such as the death of lowland, mangrove and barrier island forests and the severe contamination of soil and water supplies. The proliferation of invasive species like the Nipa palm further complicates this ecological

degradation, actively displacing native vegetation and disrupting the balance of these sensitive habitats.

The analysis highlights that the primary barrier to sustainable development is not a lack of resources, but a failure of management and regulation. Current efforts are hampered by weak policy implementation, inadequate monitoring, and a historical prioritization of industrial output over environmental integrity. To reverse this trend, the proposed Integrated Management Framework provides a necessary shift toward strategic policy formulation and multi-stakeholder engagement. By bringing together government, industry, and local communities, the IMF aims to bridge the gap between economic exploitation and ecological preservation.

Ultimately, the future of the Niger Delta depends on the immediate adoption of "result-oriented action plans". These must include the formal designation of protected areas in high-priority zones and large-scale habitat restoration projects to remediate contaminated sites. Implementing these conservation strategies is not merely an ecological necessity but a socio-economic imperative to secure the livelihoods and food security of the millions who depend on the Delta's bioresources. Through coordinated action and a commitment to environmental accountability, it is possible to transform the Niger Delta into a model of sustainable development that protects its unique biodiversity for future generations.

References

- Adekola, O., & Mitchell, G. (2011). The Niger Delta wetlands: threats to ecosystem services, their importance to dependent communities and possible management measures. *International Journal of Biodiversity Science Ecosystems Services & Management*, 7(1), 50–68. <https://doi.org/10.1080/21513732.2011.603138>
- Adeola M.O. (1992). Importance of Wild Animals and Their Parts in the Culture, Religious Festivals, and Traditional Medicine, of Nigeria. *Environmental Conservation*, 19(2):125-134. doi:10.1017/S0376892900030605
- Aghoghovwia, O. A., & Obomunu, B. (2022). Occlusion of the Niger Delta Rivers and creeks by aquatic weeds: Impact on humans and fisheries. *International Journal of Fauna and Biological Studies*; 9(2): 25-29
- Akani, G. C., & Luiselli, L. (2001). A survey of the cultural attitudes of people towards reptiles in the Niger Delta, Nigeria: implications for conservation. *Herpetological Bulletin*, 75, 19-24.
- Akani, G. C., Petrozzi, F., Ebere, N., Dendi, D., Phil-Eze, P., Nioking1, A. , Luiselli, L. (2015). Correlates of indigenous hunting techniques with wildlife trade in bushmeat markets of the Niger delta (Nigeria). *Vie et Milieu-Life and Environment*, 65(3), 169-174.
- Akani, G. C., Politano, E., & Luiselli, L. (2004). Amphibians recorded in forest swamp areas of the River Niger Delta (southeastern Nigeria), and the effects of habitat alteration from oil industry development on species richness and diversity. *Applied Herpetology*, 2, 1-22.
- Anejionu, O. C. D., Blackburn, G. A., & Whyatt, D. (2014). Detecting gas flares and estimating flaring volumes at individual flow stations using MODIS data. *Remote Sensing of Environment*, 158, 81–94. <https://doi.org/10.1016/j.rse.2014.11.018>

- Angaye, T. C., Ohimain, E. I., Siasia, E. P., Asaigbe, P. I., & Finomo, O. A. (2014). Larvicidal activities of the leaves of Niger Delta mangrove plants against *Anopheles gambiae*. *Sky Journal of Microbiology Research*, 2(7), 032-036.
- Angaye, T., & Lelei, K. E. (2025). The Impacts of Resource Exploitation in the Niger Delta: Highlighting the Urgent Need for Biodiversity Conservation. *Greener Journal of Ecology and Ecosolution*, 6(1), 1–7. <https://doi.org/10.15580/gjee.2025.1.032425046>
- Baba, S., Chan, H. T., & Aksornkoae, S. (2013). Useful products from mangrove and other coastal plants. *ISME mangrove educational book series*, 3, 45-47.
- Chidumeje, N. P. O., Kumar, L., & Taylor, S. (2015). The Niger Delta wetland ecosystem: What threatens it and why should we protect it? *African Journal of Environmental Science and Technology*, 9(5), 451–463. <https://doi.org/10.5897/ajest2014.1841>
- Chris, D. I., & Oghenetekevwe, E. (2023). Impact of Artisanal Crude Oil Refining Effluents on Interstitial Water at a Mangrove Wetland, Asari-Toru Axis of Sombreiro River, Rivers State. *International Journal of Environment and Geoinformatics*, 10(2), 12–23. <https://doi.org/10.30897/ijegeo.1132992>
- Diyaolu, O. A., Oluwabusola, E. T., Attah, A. F., Olori, E. O., Fagbemi, A. A., Preet, G., Soldatou, S., Moody, J. O., Jaspars, M., & Ebel, R. (2022). Can Crude Oil Exploration Influence the Phytochemicals and Bioactivity of Medicinal Plants? A Case of Nigerian *Vernonia amygdalina* and *Ocimum gratissimum*. *Molecules*, 27(23), 8372. <https://doi.org/10.3390/molecules27238372>
- Egonmwan, R. I. (2007). Thermal Tolerance and Evaporative Water Loss of the Mangrove Prosobranch *Tympanotonus fuscatus* var. *radula* L. (Cerithiacea: Potamididae). *Pakistan Journal of Biological Sciences*, 10: 163-166. **DOI:** [10.3923/pjbs.2007.163.166](https://doi.org/10.3923/pjbs.2007.163.166)
- Egonmwan, R. I. (2008). The Ecology and Habits of *Tympanotonus fuscatus* var. *radula* (L.) (Prosobranchia: Potamididae). *Journal of Biological Sciences*, 8: 186-190. **DOI:** [10.3923/jbs.2008.186.190](https://doi.org/10.3923/jbs.2008.186.190)
- Ewim, D. R. E., Orikpote, O. F., Scott, T. O., Onyebuchi, C. N., Onukogu, A. O., Uzougbo, C. G., & Onunka, C. (2023). Survey of wastewater issues due to oil spills and pollution in the Niger Delta area of Nigeria: a secondary data analysis. *Bulletin of the National Research Centre*, 47(1). <https://doi.org/10.1186/s42269-023-01090-1>
- Head, I., Jones, D. & Larter, S. (2003). Biological activity in the deep subsurface and the origin of heavy oil. *Nature* **426**, 344–352. <https://doi.org/10.1038/nature02134>
- Igu, N. I., & Marchant, R. (2018). Floristic Composition and Diversity of Freshwater Swamp Forests in the Niger Basin of Nigeria. *Open Journal of Forestry*, 8, 567-584. <https://doi.org/10.4236/ojf.2018.84035>
- Ita, E.O. (1994). Aquatic plants and wetland wildlife resources of Nigeria. CIFA Occasional Paper. No. 21. Rome, FAO.. 52 p
- Izah, S. C. (2018). Ecosystem of the Niger Delta region of Nigeria: Potentials and Threats. *Biodiversity International Journal*, 2(4), 338–345. <https://doi.org/10.15406/bij.2018.02.00084>

- Izah, S. C., Aigberua, A. O., & Nduka, J. O. (2018). Factors affecting the population trend of biodiversity in the Niger delta region of Nigeria. *International Journal of Avian & Wildlife Biology*, 3(3). <https://doi.org/10.15406/ijawb.2018.03.00085>
- Jonathan EL, Imoni O, Chukwuemeka P, Eteh DR. (2025). Impact of oil spills on mangrove ecosystem degradation in the Niger Delta using remote sensing and machine learning. *Journal of Geography and Cartography.*; 8(2): 11707. <https://doi.org/10.24294/jgc11707>
- Luiselli, L., Amori, G., Akani, G.C. *et al.* Ecological diversity, community structure and conservation of Niger Delta mammals. *Biodivers Conserv* **24**, 2809–2830 (2015). <https://doi.org/10.1007/s10531-015-0975-8>
- Moffat, D., & Lindén, O. (1995). *Perception and reality: assessing priorities for sustainable development in the Niger River Delta*. *Ambio*, 24 (7-8), 527-538.
- Numbere, A.O. (2018) Mangrove Species Distribution and Composition, Adaptive Strategies and Ecosystem Services in the Niger River Delta, Nigeria. In: Sharma, S., Ed., *Mangrove Ecosystem Ecology and Function*, IntechOpen, London, 17. <https://doi.org/10.5772/intechopen.79028>
- Nwankwoala, H. O., & Okujagu, D. C. (2021). A review of wetlands and coastal resources of the Niger Delta: potentials, challenges and prospects. *Environment & Ecosystem Science*, 5(1), 37–46. <https://doi.org/10.26480/ees.01.2021.37.46>
- Nwobi C, Williams M, Mitchard ET (2020) Rapid Mangrove Forest Loss and Nipa Palm (*Nypa fruticans*) Expansion in the Niger Delta, 2007–2017. *Remote Sensing* 12(14): 2344.
- Nwosu, M.O. (2010). Ethnobotany and Biodiversity Conservation in the Niger Delta, Nigeria. *International Journal of Botany*, 6(3), 310-322. [doi:10.3923/ijb.2010.310.322](https://doi.org/10.3923/ijb.2010.310.322)
- Ogbeibu, A. E., & Oribhabor, B. J. (2023). The Niger Delta Mangrove Ecosystem and Its Conservation Challenges. In *IntechOpen eBooks*. IntechOpen. <https://doi.org/10.5772/intechopen.112543>
- Ohimain, E. I. 2004. Environmental impacts of dredging in the Niger Delta; options for sediment relocation that will mitigate acidification and enhance natural mangrove restoration. *Terra et Aqua*, 97: 9-19.
- Ohimain, E. I., Benka-Coker, M O. and Imoobe, T. O. T. 2005. The impacts of dredging on macrobenthic invertebrates in a tributary of the Warri River, Niger Delta. *African Journal of Aquatic Science*: 30. 49-53.
- Ohimain, E. I., Izah, S. C. and Otobotekere, D. 2014b. Selective impacts of the 2012 water floods on the vegetation and wildlife of Wilberforce Island, Nigeria. *International Journal of Environmental Monitoring and Analysis*. 2 (2): 73 - 85
- Ohimain, E. I., Otobotekere, D., Woyengitonyokopa, B. 2014a. Unsustainable exploitation of freshwater wetland turtles and tortoises in central Niger Delta. *International Journal of Environmental Monitoring and Analysis*. 2 (2): 57 – 64
- Ohimain, E.I. (2022). Benefits and Threats of Biodiversity Conservation in Stubbs Creek Forest Reserve, Nigeria. In: Izah, S. C. (ed) *Biodiversity in Africa: Potentials, Threats and Conservation*. Sustainable Development and Biodiversity, vol 29. Springer, Singapore. https://doi.org/10.1007/978-981-19-3326-4_5

- Ohimain, E.I.; Turner, R.E.; Middleton, B.A. (2026). Mangrove ecosystems: Importance, threats and opportunities for restoration. *Water*, 18, 787–800. <https://doi.org/10.3390/w18070787>.
- Onu, N. C. H. (2003). The Oil Rich Niger Delta Region: A Framework for Improved Performance of the Nigerian Regulatory Process. *AMBIO*, 32(4), 325–326. <https://doi.org/10.1579/0044-7447-32.4.325>
- Onuoha, H. & Uzodiana, D. M. (2026). Ecosystem degradation and endangered species in the Niger Delta: understanding the biodiversity crisis. *International Journal of Innovative Environmental Studies Research* 14(1):180-192. doi:10.5281/zenodo.18882033
- Onyari, E. K., Fayomi, G. U., & Jaiyeola, A. T. (2024). Unveiling the situation of water hyacinth on fresh water bodies in Nigeria and South Africa: Management, workable practices and potentials. *Case Studies in Chemical and Environmental Engineering*, 10, 100974. <https://doi.org/10.1016/j.cscee.2024.100974>
- Raimi, M. O. (2020). Governing the Environmental Impact of Dredging: Consequences for Marine Biodiversity in the Niger Delta Region of Nigeria. *Insights in Mining Science & Technology*, 2(2). <https://doi.org/10.19080/imst.2020.02.555586>
- Richard, G., Izah, S. C., & Ogwu, M. C. (2022). Implications of Artisanal Crude Oil Refining on Sustainable Food Production in the Niger Delta Region of Nigeria. *Journal of Environmental Bioremediation and Toxicology*, 5(2), 69–77. <https://doi.org/10.54987/jebat.v5i2.775>
- Sakib, S. M. N. (2021). The Impact of Oil and Gas Development on the Landscape and Surface in Nigeria. *Asian Pacific Journal of Environment and Cancer*, 4(1), 9–17. <https://doi.org/10.31557/apjec.2021.4.1.9-17>
- Ubom, R. M. (2010). Ethnobotany and Biodiversity Conservation in the Niger Delta, Nigeria. *International Journal of Botany*, 6: 310-322. DOI: [10.3923/ijb.2010.310.322](https://doi.org/10.3923/ijb.2010.310.322)
- Ude, K., David, E. R., Governor, O. E., Nwoke, E. E., Ude, H. N., & Sarah, E. P. (2023). Effect of oil spills and gas flaring on agriculture and environment: A case of crop farmers in Niger Delta Region of Nigeria (2000 – 2022). *International Journal of Biosciences (IJB)*. <https://doi.org/10.12692/ijb/22.3.30-50>
- Udotong, J. I. R., Udoudo, U. P., & Udotong, I. R. (2017). Effects of oil and gas exploration and production activities on production and management of seafood in Akwa Ibom State, Nigeria. *Journal of Environmental Chemistry and Ecotoxicology*, 9(3), 20–42. <https://doi.org/10.5897/jece2017.0405>
- Ugochukwu, C. N. C., & Ertel, J. (2012). Negative impacts of oil exploration on biodiversity management in the Niger De area of Nigeria. *Impact Assessment and Project Appraisal*, 26(2), 139–147. <https://doi.org/10.3152/146155108x316397a>
- UNDP (2016). Niger Delta Biodiversity Project. UNDP GEF PIMS no.: 2047. United Nations Development Programme (UNDP), Global Environment Facility (GEF)
- Yetunde, E. A. (2017). Marine protected area: Prospective tool for ecosystem-based fisheries management in Nigeria. *International Journal of Biodiversity and Conservation*, 9(6), 158–166. <https://doi.org/10.5897/ijbc2016.1062>